



Hypoxaemia in Nigerian Children Presenting to the Children Emergency Ward (CHEW) of a Tertiary Hospital

Okoh Boma^{1,2} and Jaja Tamunopriye^{1,2*}

¹Department of Paediatrics, University of Port Harcourt Teaching Hospital, Rivers State, Nigeria.

²Faculty of Clinical Sciences, College of Health Sciences, University of Port Harcourt, Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Author OB designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Author JT managed the analyses of the study and managed the literature searches. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJPR/2019/v2i230101

Editor(s):

(1) Dr. Mirta Noemi Mesquita Ramirez, Hospital General Pediatrico "Ninos de Acosta Nu", Avenida de la Victoria and Bacigalupo, Reducto, Paraguay.

Reviewers:

(1) Olayinka R. Ibrahim, Federal Medical Centre, Nigeria.

(2) Dharma Lindarto, Universitas Sumatra Utara, Indonesia.

(3) Jose Luis Turabian, Health Center Santa Maria de Benquerencia, Regional Health Service of Castilla la Mancha (SESCAM), Spain.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/47741>

Received 14 December 2018

Accepted 10 March 2019

Published 30 March 2019

Original Research Article

ABSTRACT

Aim: To determine the prevalence of hypoxaemia and predictors of signs of hypoxaemia in children with various disease conditions admitted into the CHEW of a tertiary health facility.

Place and Duration: Department of Paediatrics (Children Emergency Ward). Study was done from 1st February to 30th April 2015.

Methods: This was a descriptive cross sectional study of 129 children admitted into the CHEW with various disease conditions. Biodata and clinical examination was done in all patients. Oxygen saturation (SpO₂) was determined at admission using pulse oximeter for every sick child admitted. Hypoxaemia was defined as SpO₂ less than 90%.

Results: One hundred and twenty nine children were studied. Ages of subjects ranged between 0.08years and 17 years with a mean age of 3.06 ± 3.65 years. The mean age of 3.34 ± 3.97 years for males was higher than 2.70 ± 3.22 years for females. Thirty one (24%) children had hypoxaemia on admission with 20(64.5%) with respiratory diseases. Infants ($P=.004$) and children

*Corresponding author: Email: Tamunopriyej@yahoo.com;

with respiratory disease ($P=.047$) had a significantly higher prevalence of hypoxaemia among the study group.

Chest in drawing was a common feature but grunting had the best positive predictive value of more than 80% but with low sensitivity of 3.

Conclusion: Hypoxaemia is prevalent in children who are ill and need emergency care. Respiratory diseases and infants account for a major proportion of hypoxaemic children seen in emergency wards. Chest in drawing is a common feature from different studies; presence of grunting was highly predictive in this study although had low sensitivity.

Keywords: Children; hypoxaemia; emergency care.

1. INTRODUCTION

Hypoxaemia is defined as reduced oxygen content of blood specifically in arterial blood or the reduced percentage of saturation of haemoglobin with oxygen. It is an under recognized complication of most severe illnesses in neonates and children in developing countries and a common predictor of death [1,2,3].

Hypoxaemia can be determined by measuring the level of oxygen in a sample of arterial blood or by determination of oxygen saturation in the blood using the pulse oximeter. Hypoxaemia can be defined as arterial oxygen concentration of less than 75 mmHg or blood oxygen saturation of less than 90%. [1,3] Hypoxaemia is a common manifestation of severe illnesses in children and a major contributor to mortality. Several clinical signs and symptoms have been found to predict hypoxaemia in sick children with or without acute lower respiratory tract infection, this include inability to feed, fast breathing, grunting, lower chest wall in drawing, nodding and convulsion [4,5].

Hypoxaemia is known to correlate well with disease severity and occurs mainly in diseases that impair ventilation, gaseous exchange or increase oxygen demand in the body. [6] A disease of the respiratory tract such as pneumonia which accounts for more than 2million deaths in children worldwide is commonly complicated by hypoxaemia. [6] The prevalence of hypoxaemia in children varies with disease condition and severity of illness. Prevalence of hypoxaemia in ill children range from 11 to 52% and can be as high as 73 % in children with acute lower respiratory tract illnesses. [3, 5] In a study on hypoxaemia as a measure of disease severity in young hospitalized Nigerian children with pneumonia, 41.5% had hypoxaemia with hypoxaemic children 48 times more likely to die [2].

The blood gas analysis is the gold standard for detecting hypoxemia. Other methods include pulse oximetry and less objectively clinical signs. [3] The use of the pulse oximeter is a reliable, safe, non-invasive and reproducible tool which compares well with the results from the blood gas analysis [7,8]. Its use ensures early detection and commencement of efficient treatment of hypoxaemia in sick children in resource limited setting.

The objective of this study is to determine the prevalence of hypoxaemia and predictors of clinical signs of hypoxaemia in children with various disease conditions admitted into the CHEW of a tertiary health facility.

2. METHODS

The study was a descriptive cross sectional study conducted at the Paediatric Emergency Ward of the University of Port Harcourt Teaching Hospital, Nigeria. The study population comprised of all children presenting to the Paediatric Emergency Ward, from 1st February to 30th April 2015. A convenient sample of 129 children aged 1months to 18years was studied. Exclusion criteria were children less than 1 month or 18 years or more, those SpO2 could not be determined at presentation before commencement of treatment and those whose parents did not give consent. A structured questionnaire was used to obtain information on all children studied. Children needing admission into the CHEW were consecutively recruited into the study.

The University of Port Harcourt Teaching Hospital is a tertiary health care facility in Port Harcourt, Rivers State in Southern Nigeria. It is the largest health care facility in the State and offers health care to people living in the State and its environs.

The study was approved by the Ethics and Research Committee of the University of Port

Harcourt Teaching Hospital and written informed consent was obtained from all caregivers that participated. A detailed history was taken and physical examination conducted on all children admitted into the Paediatric Emergency Ward and a clinical diagnosis made. Presence or absence of symptoms and signs of respiratory distress were particularly sought for and recorded using a study proforma. Pulse oximetry was done for all patients at presentation using a pulse oximeter (Contec CMSODL) with appropriate probe size placed on the finger and peripheral capillary oxygen saturation (SpO2) was recorded while breathing room air. Recordings were taken after stabilization of the pulse oximetry reading for one minute. Hypoxaemia was defined as SpO2 of less than 90% recorded by pulse oximetry.

All subjects were treated by the children emergency ward managing team with appropriate medications and interventions based on their individual diagnoses. They were followed up to monitor the outcome of their admission.

Data was analyzed using the Epi info version 7.1.3.3 software. The Chi square and Student's t tests were used to test for statistically significant differences in proportions and means respectively. A p value of less than or equal to 0.05 was considered as statistically significant.

3. RESULTS

A total of 129 children participated in the study, Seventy two (55.8%) males and 57(44.2%) females giving a male to female ratio of 1.3:1. Ages ranged from 0.08years to 17 years with a mean age of 3.06 ± 3.65 years. The mean age of 3.34 ± 3.97 years for males was not significantly higher than 2.70 ± 3.22 years for females. (student's $t=0.98$, $df=1$, $P=.327$). The primary diagnosis in 66 (51.2%) of the children was a respiratory disease and otherwise in the remaining 63 (48.9%).

Thirty one (24%) of the children had hypoxaemia with 20(64.5%) having primary respiratory disease. The SpO2 ranged from 54% to 99% with a mean of $91.53 \pm 8.57\%$, IQR median range of 67% to 95%. The mean SpO2 for the hypoxaemic children was $78.13 \pm 6.90\%$ while the mean SpO2 for the non-hypoxaemic children was $95.78 \pm 2.55\%$ ($P<0.001$). Table 1 shows the levels of SpO2 measured among the patients.

Table 2 shows the relationship between presence of hypoxaemia and some variables among the study group. Infants and children with respiratory disease had a significantly higher prevalence of hypoxaemia among the study group.

Table 1. Levels of SpO2 measured

SpO2 (%)	Frequency (N)	Percent (%)
≥95	69	53.5
90-94	29	22.5
75-89	27	20.9
<75	4	3.1
Total	129	100.00%

Table 2. Relationship between some variables and presence of hypoxaemia

Variable		Hypoxaemia	No hypoxaemia	Total	p
Age group	Infants	19 (35.8)	34 (64.2)	53 (42.1)	0.004*
	Children	11 (15.1)	62 (84.9)	73 (57.9)	
Gender	Males	18 (25.0)	54 (75.0)	72 (55.8)	0.390
	Females	13 (22.8)	44 (77.2)	57 (44.2)	
Primary diagnosis	Respiratory disease	20 (30.3)	46 (69.7)	66 (51.2)	0.047*
	Non-respiratory disease	11 (17.5)	52 (82.5)	63 (48.8)	

*significant

Table 3. Predictive value of some clinical signs of hypoxaemia

	Hypoxaemic patients (n=31)	Non-hypoxaemic patients (n=98)	P value	Sensitivity	Specificity	Positive predictive value	Negative predictive value
Flaring	13	29	0.107	42	70	31	79
Intercostal recession	18	40	0.05	58	59	31	82
Subcostal recession	17	30	0.009	55	69	36	83
Suprasternal recession	0	1	0.76	0	99	0	76
Grunting	1	0	0.24	3	100	100	77
Wheezing	1	1	0.424	3	99	50	76
Tachypnoea	19	47	0.102	61	52	29	81

One of the 31 cases with hypoxaemia died giving a case fatality rate of 3.2%

Table 3 shows the predictive value of some clinical signs of hypoxaemia. Tachypnoea was the most sensitive (sensitivity=61%) for hypoxaemia, followed by intercostals recession (sensitivity=58%) and subcostal recession (sensitivity=55%). Grunting (specificity=100%), wheezing (specificity=99%) and suprasternal recession (specificity=99%) were the most specific for hypoxaemia but no child had cyanosis. Only grunting had an acceptable positive predictive value of >80% in this study but however had a low sensitivity.

4. DISCUSSION

The prevalence of hypoxaemia in this study was similar to that of 20.6% in a study in Ife, Nigeria [9] and 23.8% in Kanpur, India. [10] It is however higher than the 5.8%, 11.9% and 13% found in studies done in The Gambia, [11] Chandigarh, India [12] and Enugu, Nigeria [13] respectively, and lower than 41.5% and 73% seen in Ilorin, Nigeria² and Papua New Guinea respectively. [3] Various factors including altitude, health care settings, diagnoses in subjects, age group of subjects, cut off values for hypoxaemia could have contributed to the differences seen among the various studies. Differences may also be due to prevalence of respiratory illnesses in cohort of various studies.

The significantly higher prevalence of hypoxaemia among infants (35.8%) compared to children above one year in this study is corroborated by similar studies among children in Enugu, [13] Ife, [9] The Gambia [11] and India. [12] This may be due to the fact that infants have a lower tidal volume and relative inefficient compensatory mechanisms to improve ventilation. Infants are also less unable to compensate for ventilation perfusion mismatch in

situations of increased dead space. [14] Emodi et al [13] showed an equal occurrence of hypoxia in both genders in their study as was also seen in the present study.

Hypoxaemia was significantly higher among patients with a respiratory disease compared to other diseases in this study. This is similar to findings in other studies. [2,3,11,12] While this was the finding in studies [3,11] among ill children with respiratory and non-respiratory illnesses, studies [2,12] among children with respiratory illnesses all revealed a higher prevalence of hypoxaemia in children with severe pneumonia. Pneumonia results in airway obstruction from swelling, abnormal secretions, and cellular debris. Atelectasis, interstitial edema, and ventilation-perfusion mismatch causing significant hypoxemia often accompany airway obstruction [15].

The present study showed tachypnoea and chest wall retractions (intercostals and subcostal) to be the most sensitive clinical features of hypoxaemia, and grunting, wheezing and suprasternal recession to be the most specific clinical features of hypoxaemia. In a study by Rao, et al [10] the sensitivity for hypoxaemia was highest with chest wall retraction which is similar to finding in this study. The authors also reported flaring of alar nasi, inability to feed as other strong indicators of hypoxaemia with sensitivity of 84% and 81% respectively especially in children with pneumonia. Chest in drawing was also significantly associated with presence of hypoxaemia in the study by Kuti and colleagues in Ile Ife, Nigeria. [9] In this study, grunting and wheezing had the highest positive predictive value in occurrence of hypoxaemia, also a study in Ibadan, Nigeria by Adebola and colleagues [16] showed that flaring and chest in drawing

were predictive of hypoxaemia. The use of clinical signs in determination of hypoxaemia in ill children has met diverse views. A systematic review and meta-analysis of prospective diagnostic studies that evaluated the accuracy of individual or combined clinical symptoms and signs in predicting hypoxemia among children aged <5 years with ARI, revealed that cyanosis, inability to feed, head nodding, respiratory rate >70/min and unresponsiveness/impaired arousability had high specificity but low sensitivity. [17] This report was supported by Lodha et al. [18] and Dyke et al. [19] who concluded that clinical symptoms and signs alone or in combination do not have sufficient sensitivity and specificity to predict hypoxemia in children with ALRI. Using clinical symptoms, about 20% of hypoxemic children would be missed, and 17–50% of children given supplemental oxygen would not need it in high sensitivity models based on clinical signs.[3] Pulse oximetry when used correctly provides a reliable bedside standard for detecting hypoxaemia even in developing countries. Pulse oximetry can correctly identify 20–30% more children with hypoxemias than using clinical signs alone and will ensure judicious and efficient use of oxygen therapy in resource limited setting where oxygen is not readily available. [2,20,21].

Although pulse oximetry remains the reliable means of determining oxygen saturation in children, use of clinical symptoms such as grunting, wheezing and chest in drawing may be very helpful in deciding presence of hypoxaemia in ill children where pulse oximeter is not available in resource limited settings.

5. CONCLUSION

Hypoxaemia is prevalent in children who are ill and need emergency care. Respiratory diseases and infants account for a major proportion of hypoxaemic children seen in emergency wards. Chest in drawing is a common feature from different studies; presence of grunting was highly predictive in this study although it had a low sensitivity.

The sample size in this study was small and will limit interpretations of finding but has given the authors an opportunity to obtain basic data and trend. Pulse oximeter is an invaluable tool in management of ill children and should be made available in Paediatric emergency care settings in developing countries to aid early detection and management of children with hypoxaemia.

CONSENT AND ETHICAL APPROVAL

The study was approved by the Ethics and Research Committee of the University of Port Harcourt Teaching Hospital and written informed consent was obtained from all caregivers that participated.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Lazzerini M, Sonogo M, Pellegrin MC. Hypoxaemia as a mortality risk factor in acute lower respiratory infection in children in low and middle income countries: Systematic Review and meta-Analysis. PLoS ONE. 2015;10(9):e0136166. DOI: 10.1016/j.pone.0136166
2. Abdulkadir MB, Ibraheem RM, Gobir AA, Johnson WBR. Hypoxaemia as a measure of disease severity in young hospitalized Nigerian Children with pneumonia: A cross-sectional study. S Afr J Child Health. 2015;9(2):53-56. DOI: 10.7196/SAJCH.901
3. Duke T, Blaschke AJ, Sialis S, Bonkowsky JL. Hypoxaemia in acute respiratory and non-respiratory illnesses in neonates and children in a developing country. Arch Dis Child. 2002;86:108-112. Available:www.archdischild.com
4. Weber MW, Usen S, Palmer A, Shabbar J, Mulholland EK. Predictors of Hypoxaemia in hospital admission with acute lower respiratory infection in a developing country. Arch Dis Child. 1997;76:310-14.
5. Christi MJ, Salam MA, Ashraf H, Farugue ASG, Bardhan PK et al Predictors and outcome of hypoxaemia in severely malnourished children under-fives with pneumonia: A case control design. PLoS ONE. 2013;8(1):e51376. DOI:10.1371/journal.pone.0051376
6. Subhi R, Adamson M, Campbell H, Weber M, Smith K, et al. The prevalence of hypoxaemia among ill children in developing countries: A systematic review. Lancet Infect Dis. 2009;9(4):219-27. DOI: 10.1016/S1473-3099(09)70071-4
7. Hassan A. Non-invasive monitoring of blood gases. In: Hassan A, ed. Handbook of blood Gas /Acid Base interpretation. London: Springer –Verlag. 2009;63-95.

8. Stein RT, Marostica PJC. Community – acquired pneumonia: A review and recent advances. *Pediatr Pulmonol.* 2007;42(12): 1095-1103.
DOI: org/10.1002/ppul.20652
9. Kuti BP, Adetola HH, Aladekomo TA, Kuti DK. Prevalence and predictors of hypoxaemia in hospitalized children at the emergency unit of a resource constrained centre. *Niger J Paediatr.* 2016;43(4):273 – 280.
10. Rao YK, Midha T, Kumar P, Tripathi VN, Rai OP. Clinical predictors of hypoxemia in Indian children with acute respiratory tract infection presenting to pediatric emergency department. *World J Pediatr.* 2012;8(3): 247-251.
11. Junge S, Palmer A, Greenwood BM, Mulholland EK, Weber MW. The spectrum of hypoxaemia in children admitted to hospital in The Gambia, West Africa. *Tropical Medicine and International Health.* 2006;11(3):367–372.
12. Singhi S, Deep A, Kaur H. Prevalence and predictors of hypoxemia in acute respiratory infections presenting to pediatric emergency department. *Indian J Crit Care Med [serial online]* 2003 [cited 2014;7:118-23.
Available:<http://www.ijccm.org/text.asp?2003/7/2/118/5739>.
13. Emodi IJ, Ikefuna AN, Ubesie AC, Chukwu BF, Chinawa JM. Assessment of clinical features and haematocrit levels in detection of hypoxaemia in sick children. *African Journal of Respiratory Medicine.* 2011;7(1):11-13.
14. Qureshi SA. Measurement of respiratory function: An update on gas exchange. *Anaesthesia & Intensive Care Medicine.* 2011;12:490– 5.
15. Sandora TJ, Sectish TC. Community-Acquired Pneumonia. In: Kliegman RM, Behrman RE, Jenson HB, Stanton BF editors. *Nelson Textbook of Paediatrics*, 19th ed. Philadelphia. Saunders Elsevier; 2011.
16. Adebola E, Orimadegun Babatunde O, Ogunbosi Shannon S, Carson. Prevalence and predictors of hypoxaemia in respiratory and non respiratory primary diagnosis among emergently ill children at a tertiary hospital in South Western Nigeria. *Royal Society of Tropical Medicine and Hygiene.* 2013;107(11):699-705.
17. Zhang L, Mendoza-Sassi R, Santos JC, Lau J. Accuracy of symptoms and signs in predicting hypoxaemia among young children with acute respiratory infection: A meta-analysis. *Int J Tuberc Lung Dis.* 2011;15:317–25. [PubMed].
18. Lodha R, Bhadauria PS, Kuttikat AV, Puranik M, Gupta S, Pandey RM, et al. Can clinical symptoms or signs accurately predict hypoxemia in children with acute lower respiratory tract infections? *Indian Pediatr.* 2004;41:129–35.[PubMed].
19. Dyke T, Lewis D, Heegaard W, Manary M, Flew S, Rudeen K. Predicting hypoxia in children with acute lower respiratory infection: A study in the highlands of Papua New Guinea. *J Trop Pediatr.* 1995; 41:196–201.[PubMed].
20. Graham SM, English M, Hazir T, Enarson P, Duke T. Challenges to improving case management of childhood pneumonia at health facilities in resource-limited settings. *Bull World Health Organ.* 2008;86:349–55.
21. Lee WW, Mayberry K, Crapo R, Jensen RL. The accuracy of pulse oximetry in the emergency department. *Am J Emerg Med.* 2000;18:427–31. [PubMed].

© 2019 Boma and Tamunopriye; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<http://www.sdiarticle3.com/review-history/47741>